

The Potts frustrated model: relations with glasses

Giancarlo Franzese^{1,3} * and Antonio Coniglio^{2,3}

¹ *Dip. di Fisica “E.Amaldi”, Università Roma Tre, via della Vasca Navale 84, 00146 Roma, Italy*

² *Dip. di Scienze Fisiche, Università di Napoli, Mostra d'Oltremare Pad.19, 80125 Napoli Italy*

³ *Istituto Nazionale per la Fisica della Materia, UdR Napoli*

Abstract

Similarities between fragile glasses and spin glasses (SG) suggest the study of frustrated spin model to understand the complex dynamics of glasses above the glass transition. We consider a frustrated spin model with Ising spins and s -state Potts spins both with and without disorder. We study the two models by Monte Carlo simulations in two dimensions. The Potts spins mimic orientational degrees of freedom and the coupled frustrated Ising spins take into account for frustrating effects like the geometrical hindrance. We show that in this model dynamical transitions and crossovers are related to static transitions. In particular, when disorder is present, as predicted and verified in SG, a dynamical transition between high-temperatures exponential to low-temperatures non-exponential correlation functions numerically coincides with the ordering temperature of the ferromagnetic regions, i.e. the Griffiths temperature $T_c(s)$, while a crossover between power law growth of correlation times to Arrhenius law occurs near a Potts transition at $T_p(s) < T_c(s)$. In the model without disorder, where $T_c(s)$ is not defined, both dynamical transition and crossover occur at $T_p(s)$. Furthermore the static susceptibility and the autocorrelation times of quantities depending on Potts spins diverge at $T_p(s)$. This is reminiscent of recent experimental results on glass-forming liquids.

I. INTRODUCTION

Glassy systems, like supercooled liquids, polymers, granular material, vortex glasses, ionic conductors, colloids, plastic crystal and spin glasses, can be defined as materials characterized by a slowing down of one or more degrees of freedom which prevents the system to reach the equilibrium as the temperature decreases (Angell 1984, Götze 1989, Angell 1995, Ediger *et al.* 1996).

In particular, spin glasses (SG) are dilute alloys of magnetic atoms in low concentration in a non-magnetic matrix where the relevant degrees of freedom are the impurities spins interacting via random magnetic couplings. The randomness gives rise to *frustration*, i.e.

*Present address: Center for Polymer Studies, Department of Physics, Boston University, 590 Commonwealth Avenue, Boston, MA 02215

a competition that prevents the systems to minimize the global free energy (Mezard *et al.* 1987).

One of the characteristic phenomenon of glassy systems is their complex dynamics well above the calorimetric glass transition at temperature T_g . For example there is a dynamical transition at $T^* > T_g$ from high-temperatures exponential to low-temperatures non-exponential correlation functions. Another feature is the crossover at T_1 with $T_g < T_1 < T^*$ between a high-temperature power law growth of correlation times to an Arrhenius behaviour $\tau \sim \exp(\Delta E/k_B T)$, where ΔE is the activation energy. This phenomenology is well seen in experiments (Angell 1984, Angell 1995, Ediger *et al.* 1996 Andreozzi *et al.* 1998).

Here we show the numerical results on frustrated models with and without disorder, the Potts SG (PSG) and the Potts fully frustrated (PFF) model respectively, where the dynamical anomalies correspond to static transitions, giving some insights on the relation between dynamics and statics in glassy systems and clarifying the role played by frustration and by disorder. The model could be also useful to understand some recent experimental results on finite-temperature diverging static susceptibility in glassy systems (see for example Dixon *et al.* 1990, Menon and Nagel 1995, Leheny and Nagel 1997).

II. THE PSG AND PFF MODELS

Let us consider the Hamiltonian

$$H = -sJ \sum_{\langle i,j \rangle} [\delta_{\sigma_i \sigma_j} (\epsilon_{i,j} S_i S_j + 1) - 2] \quad (1)$$

where to each lattice site is associated an Ising spin $S_i = \pm 1$ and a s -state Potts spin $\sigma_i = 1, \dots, s$. The sum is extended over all nearest neighbor (NN) sites, $\epsilon_{i,j} = \pm 1$ is the sign of interaction and J is the strength of interaction. When the $\epsilon_{i,j}$ are quenched random variables we will refer to the model as the PSG, while when they are distributed in a deterministic way to frustrate each lattice cell we will refer to it as the PFF. For $s = 1$ the PSG (PFF) model recovers the Ising SG (FF) model.

The model can be interpreted as a system with s -states orientational degrees of freedom frustrated by means of coupled Ising spins which take into account for effects due to geometrical hindrance (e.g. like in plastic crystals, where the centers of mass of the molecules form a regular crystal but the molecules are frustrated respect to the orientational degrees of freedom, or in o-terphenyl or in glycerol). The way in which the frustration is introduced distinguishes this model by other models, e.g. the Potts glass (Kirkpatrick and Wolynes 1987, Thirumalai and Kirkpatrick 1998), used to study structural glasses.

Both PSG and PFF models can be mapped on a Fortuin-Kasteleyn (FK) percolation model (Fortuin and Kasteleyn 1972, Coniglio and Klein 1980), defining the FK clusters as maximal sets of spins connected by bonds activated between NN spins with probability $p = 1 - e^{-2sJ/k_B T}$ (k_B is the Boltzmann constant) when both the NN Potts and Ising spins minimize the coupling energy and using the relation

$$Z\{\epsilon_{i,j}\} = \sum_{\{S_i, \sigma_i\}} e^{-H/k_B T} = \sum_C W_s(C) \quad (2)$$

where $W_s(C) = 0$ if the cluster configuration C contains any frustrated loop (defined below), otherwise $W_s(C) = p^{|C|}(1-p)^{|A|}(2s)^{N(C)}$ where $N(C)$ is the number of clusters in the configuration C , $|C|$ is the number of activated bonds and $|C| + |A|$ is total number of interactions. A frustrated loop is a closed path that has an odd number of antiferromagnetic interactions (Coniglio *et al.* 1991). While the Hamiltonian (1) is defined only for integer $s > 1$, the partition function (2) is meaningful for any values of s (Fierro *et al.* 1999).

III. STATICS AND DYNAMICS OF THE MODELS

The numerical phase diagrams of PSG and PFF model are shown in (Franzese and Coniglio 1998, Franzese 1999) and are qualitatively reproduced in figure 1 for the 3D case. Two transition temperatures are seen. The lower temperature $T_{SG}(s)$ (or $T_{FF}(s)$) corresponds to a SG (or FF respectively) transition in the universality class of $\pm J$ Ising SG (or Ising FF) model. The higher $T_p(s)$ corresponds to the percolation transition of FK clusters and for any integer $s > 1$ marks a real thermodynamic transition in the universality class of a ferromagnetic s -state Potts model (Wu 1982). At $T_p(s)$ the susceptibility of Potts spins diverges (Franzese and Coniglio 1998, Franzese 1999). In the PSG model, above $T_p(s)$, there is the Griffiths temperature $T_c(s)$ defined as the critical temperature of a ferromagnetic model with the same number of states of the disordered model ($2s$ for the s -state PSG model). This transition is consequence of the presence of ferromagnetic regions, due to disorder, and vanishes for vanishing external field (Griffiths 1969).

From the dynamical point of view we have studied (Franzese and Coniglio 1999, Franzese 1999) for $s = 2$ both for PSG and PFF the time dependent nonlinear susceptibility

$$\chi_{SG}(t) = \frac{1}{N} \left\langle \left[\sum_i S_i(t+t_0) S_i(t_0) \right]^2 \right\rangle \quad (3)$$

(where N is the total number of spins and t_0 is the equilibrium time) that converges asymptotically for $t \rightarrow \infty$ to the usual static nonlinear susceptibility. The normalized correlation function is

$$f_x = \frac{\chi_{SG}(t) - \chi_{SG}(t = \infty)}{\chi_{SG}(0) - \chi_{SG}(t = \infty)} \quad (4)$$

with $\chi_{SG}(0) = N$. Following (Campbell and Bernardi 1994) the infinite size behavior of $f(t)$ has been extrapolated at every t plotting the data for finite linear sizes $L = 20, 25, 30, 40$ vs $1/L$.

To test the form of $f(t)$ we fitted the data *i)* with a simple exponential, finding good fits only asymptotically for long time and for high temperatures, *ii)* with a stretched exponential function $f_0 \cdot \exp[(t/\tau)^\beta]$, finding that it fails to fit the data only for short times, *iii)* with the form $f_0 \cdot t^{-x} \exp[(t/\tau)^\beta]$ suggested by Ogielski (Ogielski 1985), finding that it fits very well the data over all the time's range and the temperature's range (Franzese and Coniglio 1999).

The parameters β , τ used in the fit *ii)* and *iii)* for χ_{SG} are plotted in figures 2, 3 for PSG and PFF respectively. In the same figures is shown also the integral correlation time defined as

$$\tau_{int} = \lim_{t_{max} \rightarrow \infty} \frac{1}{2} + \sum_{t=0}^{t_{max}} f(t) \quad (5)$$

where f is the generic correlation function.

In figure 2 for PSG it is possible to distinguish three dynamical regions. A high temperatures region for $T > T^* \simeq T_c(s)$ where the correlations functions are exponential ($\beta = 1$); an intermediate region for $T^* > T > T_1 \simeq T_p(s)$ where $\beta < 1$ and the correlation time's growth is a power law forecasted by the Mode Coupling Theory (MCT) (Götze 1989) or mean field (MF) theory (see for example Bouchaud *et al.* 1998 and references therein); a lower temperatures region for $T < T_1$ where activated processes dominate the dynamics (i.e. the correlation times follow an Arrhenius law) as in the experimental glasses (Angell 1984, Angell 1995, Ediger *et al.* 1996, Andreozzi *et al.* 1998). By the way, these results show that the prediction $T^* = T_c$ given for the SG (Randeria *et al.* 1986, Cesi *et al.* 1997) is valid also in this model.

In figure 3 for PFF, where $T_c(s)$ is not defined for the absence of disorder, the intermediate region disappears and both the dynamical transition at T^* and the crossover at T_1 occur at a temperature numerically consistent with $T_p(s)$.

For the PFF model we have calculated also the correlation functions for the Potts order parameter $M = [s \max_i(M_i) - 1]/(s - 1)$ (where M_i is the density of Potts spins in the i -th state) defined as

$$f_M(t) = \left[\frac{\langle \delta(t + t_0) \delta(t_0) \rangle}{\langle \delta(t_0)^2 \rangle} \right], \quad (6)$$

(where $\delta(t) = M(t) - \langle M \rangle$ and t_0 is the equilibration time). At the Potts transition temperature $T_p(s)$ the static susceptibility related to M diverges (Franzese 1999) and the parameter β and τ for the fit forms *ii*) and *iii*) of f_M becomes less than one and diverging respectively, as shown in figure 4. The divergence of τ comes out from the observation that $\tau(T, L)$ grows for increasing L at any T and there is a cusp at a temperature consistent to $T_p(s)$ within the numerical error.

IV. CONCLUSIONS

PSG and PFF model reproduce the experimental phenomenology of fragile glasses above the glass transition, recovering the dynamical transition between exponential and non-exponential correlation functions and the crossover between the power law growth, predicted by MF theories and MCT, and the activated Arrhenius regime for the correlation times. In particular, the disordered PSG model shows three different dynamical regimes experimentally seen for spin 1/2 probes in glass-forming liquids like o-terphenyl (Andreozzi *et al.* 1998).

The models show that these dynamical transitions and crossovers can be related to static thermodynamic transitions: in the PSG to the Griffiths transition and to the Potts transition. while in the PFF model only to the Potts transition. The mapping of these models on FK percolation model, show that the Potts transition coincides with a percolation transition where frustrated loop are excluded, giving a geometrical interpretation of the effects of frustration.

Furthermore at the Potts transition the susceptibility associated to the Potts variables diverges. This result is reminiscent of recent experimental findings (see for example Dixon *et al.* 1990, Menon and Nagel 1995, Leheny and Nagel 1997). In the models presented here this divergence is explained by the ordering of the orientational degrees of freedom. If such a divergence is present also in structural glasses like plastic crystals is matter of debate (Schneider *et al.* 1999).

Moreover the PFF model emphasizes that dynamical anomalies are not related to the presence of frustration *and* disorder, but also to the sole frustration and in this case T_p , that is the ordering temperature of the orientational degrees of freedom, marks the onset of these anomalies. This could be relevant in the study of dynamical behavior of experimental FF systems like Josephson junction arrays.

- Andreozzi, L., Giordano, M., and Leporini, D., 1998, *J. Non-Cryst. Solids*, **235-237**, 219.
- Angell, C.A, 1984, *Relaxation in Complex Systems*, edited by Ngai, K.L, and Wright, G.B. (Washington: Office of Naval Research).
- Angell, C.A, 1995, *Science*, **267**, 1924.
- Bouchaud, J.P., Cugliandolo, L.F., Kurchan, J., Mezard, M., 1998. *Spin Glasses and Random Fields*, edited by Young, A.P., (Singapore: World Scientific).
- Campbell, I.A., and Bernardi, L., 1994, *Phys. Rev. B*, **50**, 12 643.
- Cesi, F., Maes, M., and Martinelli, F., 1997, *Commun. Math. Phys.*, **188**, 135.
- Coniglio, A., di Liberto, F., Monroy, G., and Peruggi, F. 1991, *Phys. Rev. B*, **44**, 12 605.
- Coniglio, A., and Klein, W., 1980, *J. Phys. A*, **13**, 2775.
- Dixon, P.K., Wu, L., Nagel, S.R., Williams, and J.P. Carini, 1990, *Phys. Rev. Lett.*, **65**, 1108.
- Dixon, P.K., Wu, L., Nagel, S.R., Williams, and J.P. Carini, 1991, *Phys. Rev. Lett.*, **66**, 960.
- Ediger, M.D, Angell, C.A, and Nagel, S.R., 1996, *J. Phys. Chem*, **100**, 13 200.
- Fierro, A., Franzese, G., de Candia, A., and Coniglio, A., 1999, *Phys. Rev E*, **59**, 60.
- Fortuin, C.M., and Kasteleyn, P.W., 1972, *Physica*, **57**, 536.
- Franzese, G., 1999, preprint.
- Franzese, G., and Coniglio, A., 1998, *Phys. Rev. E*, **58**, 2753.
- Franzese, G., and Coniglio, A., 1999, preprint <http://xxx.lanl.gov/abs/cond-mat/9803043> in press on *Phys. Rev. E*, **59**, vol. 6.
- Götze, W., 1989, *Liquids, freezing and glass transition*, edited by Hansen, J.P., Levesque, D., Zinn-Justin, J. (Amsterdam: North Holland).
- Griffiths, R.B., 1969, *Phys. Rev. Lett.*, **23**, 17.
- Kirkpatrick, T.R., and Wolynes, P.G., 1987, *Phys. Rev. B*, **35**, 3072.
- Leheny, R., and Nagel, S.R., 1997, *Europhys. Lett.*, **39**, 447.
- Menon, N., and Nagel, S.R, 1995, *Phys. Rev. Lett.*, **74**, 1230.
- Mezard, M., Parisi, G., and Virasoro, M.A., 1987, *Spin Glass Theory and Beyond* (Singapore: World Scientific).
- Ogielski, A.T., 1985, *Phys. Rev. B*, **32**, 7384.
- Randeria, M., Sethna, J.P., and Palmer, R.G., 1986, *Phys. Rev. Lett.*, **57**, 245.

Schneider, U., Brand, R., P. Lunkenheimer, and A. Lloidl, 1999, preprint <http://xxx.lanl.gov/abs/cond-mat/9902318>.
Thirumalai, D., and Kirkpatrick, T.R., 1998, *Phys. Rev. B*, **38**, 4881.
Wu, F., 1982, *Rev. Mod. Phys.*, **54**, 235.

FIGURES

FIG. 1. Qualitatively phase diagram for the PSG model in $3D$ as function of s . Solid lines mark real thermodynamic phase transitions, while dotted line marks the vanishing Griffiths transition. Cross marks Potts vanishing transition in the $s = 1$ ($\pm J$ Ising SG) case. An analogous phase diagram holds for the PFF model with a FF Ising phase at the place of the SG phase and without the G phase.

FIG. 2. PSG for $s = 2$: β and τ parameters used to fit the nonlinear susceptibility correlation function for $L \rightarrow \infty$ (from Ref.[breve]). Circles are the parameters for the form *iii*), triangles for the form *ii*); squares are the integral correlation times. Where not shown the errors are smaller then symbols size. Arrows show $k_B T_c/J = 3.641$ and $k_B T_p/J = 2.925 \pm 0.075$.

FIG. 3. PFF with $s = 2$: As in the previous figure. Arrow shows $k_B T_p/J = 2.76 \pm 0.05$.

FIG. 4. PFF with $s = 2$: as in the previous figure but for the Potts order parameter's correlation function f_M .







